



ለፕላንቢት ምድር

ᠳᠤᠵᠢ ᠴᠡᠭᠦᠨᠠᠩᠲᠤᠨᠣᠯᠪᠤᠰᠤᠨ: Wednesday, April 23, 2025

Period of operation: from 2025-04-13 to 2025-09-15

Tamzin Blewett
University of Alberta
118 and 85th ave
Edmonton Alberta T6G 2R3
Canada
Tel: 780.492.9057, Email: tblewett@ualberta.ca

ፍጹሙ ለሥራ ለሥራ ለሥራ ለሥራ ለሥራ

Inuinnaqtun: Hilaqyuaqmi hilap allanguqnia hatqiqtai ayyikkutaittut ihuinit tapkuat qangani ataanilu (naunaqkutariplogu, Ukiurtaqtuq tamnal Ukiurtaqtuq kayumiktumik hutqiknia uunaqpalliqninut nalaanit. Piplugit, qiqumainnaqnit Ukiurtaqtuq avikhimani qahuvalliqitut, tariuplu hikua hikhimani ikiklivalliqitut, pipkaqhugu umiaqpait huliniit, uqhuqyuaq kaasitlu amuyauni, havikhat uyarakhiuqni, pulaaraqutlu, tamaqmik pilaqninut hatqiqni halumailrut ilavalliqninut umiaqpait aulani umiaqpaitlu aulani. Iqaluktutiak tamnal Tulakvik Iqaluit piniqhamik piyaulat ilagiqni haviit halumailrut ingattaqyumininut havikhat amuyuani tahapkunani inaitni uqhuqyuanutlu halumaiqnit avatainut. Hulivalliq, atuqni pukkittuq sulfur uqhuqyuat atuqpaliqtauplutik Ukiurtaqtumi mikhigiqahugit hilaqut halumaiqnit tahapkuat ilagiqni huliniit. Kihimik, tahapkuat nutangutqiyat uqhuqyuat pilgit allatqinik avugiyauvaktut piqaqnit, tapkuat aktuanit qayurnaqnigmi atuliqnit Ukiurtaqtuq piqarutai kangiqhimayauttiangitni. Ilagiplugitlu, tatya imaq nakuunia hivulirutai piplugu ikpigilaqnit angutikhat uunaqniuvalliqninut nalaanitnit.

Piplugu tamna, tahapkuat hivulirutit pihimaitai tapkuat taihimayut avatiliqutit pityutai piqatai niglamayuni imaq avatait uumatyutitlu, pingitailu tapkunanga ikpigilaqni Ukiurtaqtuq uumayut halumailrut. Taimaittumik, nakuuhivallirumayavut havariyaqtaqta hapumminit Ukiurtaqtuq uumatyutai tapkunanga inungnit piqatai ulapihautit. Una naunaiyaut piniaqhimayuq naunaiqni piyakhai uvaluniit pingittaq Ukiurtaqtuq uumayut ikpigilaqniqhauni halumaittunut tapkuat uunaqni uumayut tatyatuqtayut ihuaqhini imaq nakuunia atuqpaknit. Taimailuqhugu, niriuktugut pivaliatitni nakuutqiyamik hivulirutai taihimayut hapumminia Ukiurtaqtuq avatai. Una havanguyuuq piniaqhimayuq kangiqhini Ukiurtaqtuq uumayuvalui ikpigilaqni malruk qanurittuni tuqunait: haviit halumailrut (copper, zinc, cobalt, nickel, tamnalu lithium) tamnalu uqhiqyuuaq turangani avugiyauvaktut (ilautitlugit perfluoroalkyl tamnalu polyfluoroalkyl hunaunit (PFAS) ahiilu hunat talvanga pukittuq sulfur uqhuquuat). Hatqiqniaqtavut pingahut qanurittuni uumayuvaluit nalvauyut Ukiurtaqtumi tahapkuat tuqunait piyauyumayut atuqhugu Ukiurtaqtuq turangayut katiqhuqnit qanuritnit (naunaipkutariplugu, uunaqni, imaq avigiyauttaqni). Uukturaqhugit allatqit katiqhuqnit hatqiqni, pinahuaqtavut naunaiqni amihuniqhat qanurililat uumayut attaqtuhigiqnulu ilihimanivut ikayuqni nunaluuyut, piqatauyut, maligaliuqtut hapummini Ukiurtaqtuq avatigiyai. Tariup hilkua aqayait, uugavuluit, tapkuatlu Daphnia (imaqmi kumavauit) katitauniat talvani avataanilu Ikaluktutiak, NT. Taiplugu, aqayait uugavuluitlu katitauniat talvani Qiqaiyaluarvia 2025 talvanga tariup hikua akungani Ikaluktutiak tamnalu Dease Strait tariunga, pititlugu Daphnia imaqilu naunaiyaqni katitauniat talvani Niqiliqivik talvanga Ikaluktutiak (Greiner Tahiq). Tapkuat hatqiqni talvani Qiqaiyaluarvia atukhaqniat pingahunik havaguhit, tapkuatlu Niqiliqivik atuqniat tallimat havaguhit. Tamaita uuktugayut piniait hivuranaittumik hiamaktailini ilauni Kanatamiuni Quttiktuq Ukiuqtartuq Naunaiyivik (CHARS) naunaiyavit, imaqilu atuqnia uukturaqni halumaqhagauniat iqaqniahaqlugit ahivaqni kitutliqak hivuranaqtut hunat. Allatqit halumaqtiqni pityuhit, tapkuatut uumani iqalukpit halumaqhautit, haviknut nipittaqni, huliplituklu paulgit, atuqniaqni ahivaqni havit avigiyauyutlu, atuqpiqhugit atulaitni hivuranaqtut halumailrut hatqiumani avatigiyauyunut atuqtitlugu naunaiyainit havanguyuuq. Kinguagut hatqiqnit iniqtiqniagut, tamna uumayuvaluit aulaqtitauniat talvunga Quebec City (Institut national de la recherche scientifique; INRS) tamnalu/uvaluniit Edmonton (Universitynga Alberta) hulivalli q naunaiyautit takuni qanuq tahapkuat halumailrut piruqhaqtut akuniyuuq aktuanilu Ukiurtaqtuq uumayuvaluit amihuvalli qni uiguni pivikhiat havagutai. Una havakvikni ilagiya 3 ukiut NSERC Katutyiqatai manikhat tamnalu 3 ukiut DFO-kut manikhat; atuqhugu havaqatigini manikhat, nunalikni piqatigikni atuqniat tuhaumatyutit naunaiyainivut piqatigini nunaluuyut Ikaluktutiak. Tamna havanguyuuq havaqatigit havariniaqta nunalikni katimanit hatqiqtaunit ukiuq tamaat havanguyumut. Tahapkuat huliniit ilaqaqiat takukhautitni hanalrutit atuqni uuktutini naunaiyautit tapkualuttauq nutanguqnit naunaiyainiq havanguyuuq qanuritnit tuhagakhatlu tapkuninga tatyatuq qanuritnit maligaitlu avataini atuqpiarialgit havikhait. Atuqtitlugu maniqami naunaiyautit, parnaktugut pihimani ilauyut nunalikni angutikhaliqiyit (HTO) piqatiginivut atuqpiariangi ikayuqtavutlu nunalikni nunaluuyuuq huli ilitni nanminiq nunalikni ilauyunut. Kingulli qpamik, parnaktugut piqatauniq nunaluuyumi niriqatigiknit piqatigiplugit nunalikni HTO-kut piruqhaqpalli qni atatyutai nunalikni nunaqaqaqtut nunaluuyut.

Personnel

Personnel on site: 8

Days on site: 65

Total Person days: 520

Operations Phase: from 2025-04-13 to 2025-09-15

Λ Γ Δ Ε Ζ Η Θ Ι Κ Λ Μ Ν Ξ Ο Π Ρ Σ Τ Υ Φ Χ Ψ Ω

ᐱᕈ	ᖃᓄᐸᑦᑐᒥᑦ ᐱᕋᕆᐸᖃᑦᓂᐸᖅ<	ᑭᔪᑦ ᓄᐳᖃᑦᑯᑦᓂ	ᑐᕗᐤᐤᙵᕐᕐ ᓄᐳᑦ < ᖃᓄᖅ ᐸᑐᓴᐤᐸᐤᖅᕈᐤᐤᕆᐸᐤᖅᓴᑦᓂᖅ	ᐸᑦᕐᓂᑦᖃᑦᐱᐤᐳᑦᐸᖅ< ᐸᓄᖅᓄᑦ ᐳᕐᖅᑲᐤᖃᑦᑲᑦᓂᑯᑦᑐ ᑲᐸᑦᕈᐤᓂᑐᖅᐤᕋᑦᑐᖅ	ᖃᓂᑦᓂᖅ<ᖅ ᓄᐳᕋᕏᙵᐤᕐᖅ ᐸᕐᐤᐤ ᕐᕗᕐᕐᕐᕈᐼᐤᐳᑦ ᕈᐳᑦ
Lake where Cambridge Bay drinking water is taken from	Sampling sites	Municipal	Site was chosen as a potential collection location for water to look at metal contaminants in the drinking water that feeds Cambridge Bay	N/A	This sampling site is ~2km North of Cambridge Bay
Outflow of sewage lagoon, leading into river/bay	Sampling sites	Municipal	Site was chosen as a potential collection location for water collection to look at metal contaminants flowing into the bay from the Cambridge Bay sewage lagoon.	N/A	Outflow of Sewage lagoon is ~1.3km Northwest of Cambridge Bay
Downstream of sewage lagoon, closest to outflow	Sampling sites	Municipal	Site was chosen as a potential collection location for water collection to look at metal contaminants just outside of the sewage lagoon outflow, in the bay outside of the town.	N/A	Downstream of Sewage lagoon, close to outflow is ~1.25km Northwest of Cambridge Bay
Downstream of sewage lagoon, intermediate distance	Sampling sites	Municipal	Site was chosen as a potential location for water collection to look at metal contaminants downstream of the sewage lagoon outflow, in the bay outside of town.	N/A	Downstream of Sewage lagoon, intermediate to outflow is ~1km Northwest of Cambridge Bay
Downstream of sewage lagoon, longest distance	Sampling sites	Municipal	Site was chosen as a potential location for water collection to look at metal contaminants farthest downstream from sewage lagoon outflow, but in close proximity to town.	N/A	Downstream of Sewage lagoon, farthest from outflow is ~0.9km Northwest of Cambridge Bay
Greiner Lake site for Daphnia collection	Sampling sites	Municipal	This lake was chosen for water flea collection because of previous work outlining the food web dynamics (Grosbois, G., M. Power, M. Evans, G. Koehler, and M. Rautio. 2022. Ecosphere 13(1):e03881. 10.1002/ecs2.3881) and sediment core analysis of ecological changes over time in this water body (Kivila, E., Rantala, M., Antoniades, D., Luoto, T., Nevalainen, L., Rautio, M. 2022. CATENA 211:105969. 10.1016/j.catena.2021.105969).	N/A	Greiner Lake is 12 km North of Cambridge Bay and 5km West of Ovayok Territorial Park

Cambridge Bay Sea-ice associated algae collection and copepod collection area	Sampling sites	Municipal	This site was chosen due to its proximity to Cambridge Bay, while still containing sea ice that can be cored for algae. Ice-algae eating copepods will also be collected under the ice.	N/A	Collection area is ~2km Southeast of Cambridge Bay and 14km Southwest of Ovayok Territorial Park
---	----------------	-----------	---	-----	--

ᓄᓇᓕᓯᓪᓗ ᐃᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ ᐃᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ

ᓄᓇᓕᓯᓪᓗ	ᐃᓕᓯᓪᓗ	ᓄᓇᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ	ᓄᓇᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ
ᐃᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ	Beverly Maksagak	Ekaluktutiak Hunters & Trappers Organization	2024-09-23
ᐃᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ	Gabriel Ferland	Viventem Science Support Agency	2024-11-18
ᐃᓕᓯᓪᓗ ᓄᓇᓕᓯᓪᓗ	David Hik	POLAR Knowledge Canada	2023-08-17

ᑕᐃᕐᕐᕐ ᐱᕐᕐᕐ ᐃᕐᕐᕐᕐᕐ

ᐃᕐᕐᕐᕐ ᐱᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐ ᐱᕐᕐᕐᕐᕐ:

ᑕᐃᕐᕐᕐ ᐱᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐ

ᑕᐃᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐ	ᐃᕐᕐᕐᕐᕐᕐᕐ ᑕᐃᕐᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐᕐ	ᐃᕐᕐ ᐃᕐᕐᕐᕐᕐᕐ	ᐃᕐᕐᕐ ᐃᕐᕐᕐᕐ/ ᐃᕐᕐᕐᕐᕐᕐᕐᕐᕐᕐ	ᐃᕐᕐᕐᕐᕐᕐ
ᐃᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐ	Applied for a Research License to conduct the proposed research around Cambridge Bay	Applied, Decision Pending		
ᐃᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐᕐᕐ	Applied for a License to Fish for Scientific Purposes, but only to have DFO approval for the research. We will not be collecting any fish or fish tissue for this year.	Applied, Decision Pending		
ᐃᕐᕐᕐ ᐃᕐᕐᕐᕐ, ᐃᕐᕐᕐᕐᕐᕐᕐᕐ	Applied for a Nunavut Wildlife Research Permit due to the collection of algae and small crustaceans (copepods and Daphnia).	Applied, Decision Pending		
ᐃᕐᕐᕐ ᐃᕐᕐᕐᕐᕐ ᐃᕐᕐᕐᕐᕐ	Applied for Approval for the use of Water or Deposit of Waste Without a License due to the low volume of water (0.024 m3/day) we are expecting to use.	Applied, Decision Pending		
Hunters and Trappers Associations/Organizations	Approved for the proposed research from the Ekaluktutiak Hunters & Trappers Organization (see letter of support)	Active	2025-04-01	2026-10-01

Project transportation types

Transportation Type	ᐃᕐᕐᕐ ᐃᕐᕐᕐᕐᕐᕐ	Length of Use
Air	Researchers will take flights from southern Canada (Edmonton, Waterloo, Quebec City) to Cambridge Bay	
Water	Researchers will travel across ice sheets outside of Cambridge Bay to collect sea ice algae & copepods, and potentially via boat to collect Daphnia on Greiner lake if necessary in August	
Land	Researchers will travel across land to Greiner Lake for collection of	

	Daphnia	
--	---------	--

Project accomodation types

◁P^L,

A^cd^c d^ar^t^b d^c^bCd^cσd^ah^t^b Δc^bh^cp^cn^cr^c ΔjCΔ^c, Γ^c-jΔp^cn^c, ^bh^cLc^j^b, qe^cp^c d^ar^t^b-j

[illegible][illegible]

						chemical will not leave the lab facility at CHARS.
Ethanol	hazardous	1	1	1	Liters	500 mL, used for sanitization of work stations and equipment in the lab at CHARS. Contained in plastic container, double bagged, stored in a fume hood cabinet with other corrosive materials, inside a plastic bin. This chemical will not leave the lab facility at CHARS.
Nitric Acid (HNO3)	hazardous	4	1	4	Liters	Acid used for dissolving metals in exposure waters and cleaning glassware in the lab at CHARS. Contained in plastic container, double bagged, stored in a fume hood cabinet with other corrosive materials, inside a plastic bin. This chemical will not leave the lab facility at CHARS.

ΔL^{cb} <D^{cb}C>L^{cb}D^{cb}

Δ ^c <L ^{cb} <D ^{cb} C>σ<L ^{cb} D ^{cb}	σ _Δ ^{cb} ΔΓ ^{cb} C ^{cb} C ^c σ<L ^{cb} D ^{cb}	ΔP ^c ΔΓ ^{cb} C ^{cb} C ^c σ<L ^{cb} D ^{cb}
0	Saltwater (May; ~50L)/freshwater (August; ~50L) collected from sampling sites via pump or bucket. Water will be brought to Canadian	May – Sea-ice associated algae collection and copepod collection area; August – Greiner lake

	High Arctic Research Station for culturing organisms.	
--	--	--

ለርብሰታዊነት ለርብሰታዊነት	ፍጥነት 4bCd ^c	ፍጥነት 4bCd ^c ካላቸው ስላሉት	ፍጥነት 4bCd ^c 4bCd ^c	ካላቸው ስላሉት
Researching	4bCd ^c 4bCd ^c	40 L	5% v/v acid bath (prepared in ultrapure water) will be neutralized with acid-neutralizing powder (Calcium carbonate), then the pH of the solution will be checked to confirm neutral pH (pH = 7). Water will then be disposed in the sink under flowing water.	No additional treatment procedures
Researching	Other, Metal-spiked freshwater	3 L per metal, 15 L total for freshwater	We will generate up to 12 L of metal-spiked freshwater with about 100-500 µg/L of Co, Cu, Zn and Ni (environmentally relevant levels). These metals will be removed from the freshwater by complexation with an ion-exchange resin (chelex type), using a batch method that has been optimized to efficiently remove metal from water since the 60s. Following this treatment, the resin will be filtered out of the water (and brought to Edmonton) while the metal-free water will be disposed down the sink at CHARS, under flowing tap water.	We will generate up to 3L of freshwater contaminated with lithium. This monovalent metal will have a relatively low affinity for the chelex resin, so the above-mentioned removal technique is not optimal. This 3 L wastewater will be shipped from CHARS to Edmonton for disposal.
Researching	Other, Metal-spiked seawater	6.6 L per metal, 33 L total for seawater	We will generate up to 26 L of metal-spiked seawater with about 100-500 µg/L of Co, Cu, Zn	We will generate up to 6.6L of seawater contaminated with lithium. This monovalent metal will have a relatively low affinity for the chelex resin, so the

			and Ni (environmentally relevant levels). These metals will be removed from the seawater by complexation with an ion-exchange resin (chelex type), using a batch method that has been optimized to efficiently remove metal from seawater since the 60s. Following this treatment, the resin will be filtered out of the water (and brought to Quebec) while the metal-free water will be disposed down the sink at CHARS, under flowing tap water.	above-mentioned removal technique is not optimal. This 6.6 L wastewater will be shipped from CHARS to Quebec for disposal.
Researching	Other, Fuel-spiked seawater	6.6 L of fuel-spiked seawater	Exposure water will be treated with activated charcoal to remove fuel, then both water and charcoal will be shipped back to the University of Alberta for disposal	No additional treatment procedures

Environmental Impacts

Environmental impacts of this study are expected to be minimal due to toxicant exposures occurring in controlled laboratory environments in small volumes of water at the Canadian High Arctic Research Station (CHARS). Experiments will utilize increasing concentrations of metals (Nickel, Copper, Zinc, Cobalt, Lithium) and low sulfur fuel oil (including oil co-contaminants such as antioxidant, surfactants and per- and -Polyfluorinated Substances (PFAS)) to identify the sensitivity of Arctic animals (sea-ice algae, copepods, and water fleas) to potential increases in aquatic concentrations. High concentrations of toxicants in exposure water will be used during this experiment and nitric acid baths used to clean glassware/plasticware between experiments. Therefore, mitigation measures will be utilized to reduce potential pollution of Nunavut waters. After exposures, water will be treated with biochar and chelex resin treatment to remove dissolved metals before disposing water down sinks under flowing water. Biochar will be shipped to Edmonton (University of Alberta) or Quebec City (INRS) for appropriate disposal. For oil exposures, water will be treated with activated charcoal to reduce concentrations and/or shipped back to Edmonton for disposal. Additionally, nitric acid bath will be neutralized with acid-neutralizing powder, then the pH of the solution will be checked to confirm neutral pH (pH = 7). Water will then be disposed in the sink under flowing water.

Additional Information

SECTION A1: Project Info

SECTION A2: Allweather Road

SECTION A3: Winter Road

SECTION B1: Project Info

SECTION B2: Exploration Activity

SECTION B3: Geosciences

SECTION B4: Drilling

SECTION B5: Stripping

SECTION B6: Underground Activity

SECTION B7: Waste Rock

SECTION B8: Stockpiles

SECTION B9: Mine Development

SECTION B10: Geology

SECTION B11: Mine

SECTION B12: Mill

SECTION C1: Pits

SECTION D1: Facility

SECTION D2: Facility Construction

SECTION D3: Facility Operation

SECTION D4: Vessel Use

SECTION E1: Offshore Survey

SECTION E2: Nearshore Survey

SECTION E3: Vessel Use

SECTION F1: Site Cleanup

SECTION G1: Well Authorization

SECTION G2: Onland Exploration

SECTION G3: Offshore Exploration

SECTION G4: Rig

SECTION H1: Vessel Use

SECTION H2: Disposal At Sea

SECTION I1: Municipal Development

ᐱᓐᓇ ᐱᓇᐱᓂᐱᓐ ᐱᓇᐱᓂᐱᓐ ᐱᓇᐱᓂᐱᓐ

The sampling for this project will take place in two different physical environments. The first will be sampling on the sea ice of the Cambridge Bay inlet area in May 2025. This sampling will occur on the ice sheet ~2km Southeast of Cambridge Bay and 14km Southwest of Ovayok Territorial Park. The exact location of our sampling will be determined when we are already in the field and based on sea ice conditions, safety, appropriateness, and wildlife observations to minimise disruption. Our local team, Viventem, will be guiding this research and helping inform where the sampling sites should be. The area of the proposed sea-ice sampling is not within any known sensitive marine habitat areas, recreational areas, or heritage sites. The second sampling area will be in Greiner Lake in August 2025. Greiner Lake is a shallow basin lake (average depth 3-5m and maximum depth of 12m) with an area of 47 km². It is located ~5km from Ovayok Territorial Park, and 12km North of Cambridge Bay. The lake is covered with 2m of ice in winter, and the open water period is 2 months between early July and mid-September, aligning with our proposed sampling period. 40% of the lake is <2m deep, allowing sampling close to the shoreline to minimise potential disruption. The sampling at Greiner Lake will not disturb any designated park areas or heritage sites. Greiner Lake is used for fishing and contains Arctic char, lake whitefish, and lake trout. Our local guide, Viventem, will ensure our zooplankton sampling does not disturb fishing or subsistence practices. There have been some archaeological sites within the area, looking at historical settlements. However, these are mainly along the Ekalluk River and outside the current municipal area and will not be near our sampling sites. By-law 37 of the Hamlet of Cambridge Bay on Noise Control means that there are minimal noise levels within the region, and all activities are required to reduce unnecessary or excessive noise disturbances. Neither sampling site is within the Valued Ecosystem Components of Nunavut, including Caribou ranges, seal hunting areas, and breeding grounds. All efforts will be made to respect and preserve all natural, cultural or historical resources.

ᐱᓐᓇ ᐱᓇᐱᓂᐱᓐ ᐱᓇᐱᓂᐱᓐ ᐱᓇᐱᓂᐱᓐ

Algal samples will be collected in May 2025 from 9cm ice cores that contain sea-ice associated algae in the bottom 5cm (mainly diatoms). Sea ice is ~2m thick and will be cored using our Kovacs Ice Corer, and as few cores as possible will be used to collect sufficient algae for exposures back at CHARS (up to 12 cores). Approximately 500 individual copepods (about 1-5mm in size) will be collected in May 2025 by vertical trawl with a plankton net. The plankton net will be lowered up to 10m through a 21cm hole in the ice. Due to the infinitely small amount of water and organisms collected in relation to the vast under-ice ecosystem, the disturbance of this ecosystem will be minimal. Greiner lake sampling will occur in August along the shoreline to collect water fleas (*Daphnia* spp., estimated to be 21% of the zooplankton community) that will be present in the water column. Water fleas will be brought back to CHARS for exposures. The lake contains a large population of sea-run Arctic char (*Salvelinus alpinus*), resident lake whitefish (*Coregonus clupeaformis*), and lake trout (*Salvelinus namaycush*). However, since we are only sampling close to the shoreline, the disturbance to these fishes will be minimal. No SARA species, wildlife, or birds will be impacted by this work.

ᐱᓐᓇ ᐱᓇᐱᓂᐱᓐ ᐱᓇᐱᓂᐱᓐ ᐱᓇᐱᓂᐱᓐ

All fieldwork to sample water and our organisms will be conducted around Cambridge Bay. All exposures will be conducted inside the laboratory facility within the Canadian High Arctic Research Station (CHARS). All test waters will be fully treated prior to safe disposal (in the case of cobalt, zinc, nickel and copper-spiked waters), or

shipped to Quebec or Alberta for disposal (in the case of fuel- and lithium-spiked waters). As far as we know, there are no archeologically or culturally significant sites in the project area or in adjacent area. We will not be disturbing the palaeontological component of surface and bedrock geology. We will not be impacting any land or resource use in the area and will be following community guidelines should any of our sampling impact these operations. Our water and animal sampling will be outside of the residential area and should have minimal impact on local or regional traffic patterns. However, should our equipment or techniques impact this infrastructure, we will consult with local authorities to mitigate these impacts. Our work will have no impact on human health due to the extensive treatment plan we have for handling our relatively small volumes of exposure waters after experiments in the CHARS laboratory are completed. Additionally, our field sampling will be led by a local team (Viventem) with in-depth knowledge of the local area to minimize negative impacts.

Miscellaneous Project Information

All exposures will be contained in low volumes at CHARS and none of the solutions containing contaminants will leave the laboratories at this facility. No exposure solutions (metal- or fuel-spiked water) will be brought into the field or onto the ice during sampling of organisms. Comprehensive Spill Prevention/Plan: Handling of dangerous chemicals will follow the University of Alberta Chemicals – General Safe Handling Standard Operating Procedure (SOP; UofA_Comprehensive_Spill_Prevention.docx). If any spills occur, we will follow the University of Alberta Spill remediation protocol (UofA_Comprehensive_Spill_Plan.docx). Emergency Response Plan: Attached field activities plan (Emergency_response_plan_FAP_Blewett_Arctic_contaminant_2025.docx) contains information for Emergency Responses and mitigation of hazards. It also contains staff training and emergency contacts. Waste Management Plan/Program: Exposure waters containing metals will be treated with Chelex resin as per the attached SOP (Waste_management_plan_chelex_resin.pdf). Exposure waters containing oil (and oil co-contaminants) will be treated with Biochar as per the attached SOP (Waste_management_plan_biochar.docx).

Designated environmental areas

Designated environmental areas: Will not be impacted by this project. Ground stability: Will not be impacted by this project. Permafrost: Will not be impacted by this project. Hydrology/limnology: Will not be impacted by this project. Water quality: All lab experiments will be contained in low water volumes within the laboratory facilities at CHARS and none of the waters spiked with contaminants will be used outside this facility. Saltwater and freshwater spiked with metals or fuel will be treated to remove contaminants as per the waste section of this application. We are a team of experienced lab scientists who will be following best waste management practices, so we anticipate no environmental impact from lab waste. Climate Conditions: Will not be impacted by this project. Eskers and other unique or fragile landscapes: Will not be impacted by this project. Surface bedrock and geology: Will not be impacted by this project. Sediment and soil quality: Transport to Greiner lake will be conducted via roads where possible and guided by our local team (Viventem) to take the most appropriate and least disruptive route. Tidal processes and bathymetry: Will not be impacted by this project. Air quality: Will not be impacted by this project. Noise levels: There is the potential for ice coring to produce noise pollution for local wildlife. Therefore, we will only be conducting minimum drilling activity (<1 h cumulatively in May 2025) to collect just enough algae/copepods for our lab experiments. Since we are working in a controlled laboratory environment, we will not need much biomass for our exposures. Cores will be taken at one time to prevent repeated disturbance and guided by our local team (Viventem) who have extensive experience collecting ice cores. All activities will work within By-law 37 of the Hamlet of Cambridge Bay on Noise Control to minimize noise disruption. Vegetation: We will be collecting samples of sea-ice algae from the bottom 5 cm of about 12 ice cores (9-cm in diameter), this is the minimum amount of algae for our exposures. Wildlife: Will not be impacted by this project. Birds: Will not be impacted by this project. Aquatic species: Greiner lake houses multiple cultural and subsistence species of fish. Therefore, to minimize any impact on their wellbeing, we will minimize time at the site and avoid disturbance of fish habitat. For this work, no fish will be collected. We will be collecting zooplankton (water flea or Daphnia) and water to culture them in, making sure to collect the minimum amount of water possible. Wildlife protected areas: Will not be impacted by this project. Archaeological and cultural historic sites: Will not be impacted by this project. Employment: Our grants contain funds to hire locals to support this researcher. Therefore, we will be hiring Viventem as well as individuals from the Ekaluktutiak HTO to act as guides and support our research. Community wellness: Will not be impacted by this project. Community infrastructure: Will not be impacted by this project. Human Health: Our research will provide the basis for investigating the potential toxicological impacts of metal exposure. Overall, we hope to develop Arctic-specific guidelines for water quality that will inform future research and policy, ultimately to improve health and well-being of local Arctic communities.

Cumulative Effects

There should be minimal cumulative impacts from this research because we are visiting different sites during

our two field campaigns. All exposures will be contained in low volumes at CHARS and none of the solutions containing contaminants will leave the laboratories at this facility. Additionally, we have extensive treatment protocols for removing all contaminants from our metal- or fuel-spiked water before disposal. This means that minimal environmental repercussions will occur and pollution from our exposures will be mitigated.

Impacts

$\mathbb{A}^{\mathfrak{b}} \mathbb{C} \triangleright \sigma^{\mathfrak{a}} \tau^{\mathfrak{c}} \triangleleft \mathfrak{A} \mathfrak{B} \mathfrak{C} \triangleright \mathfrak{C}^{\mathfrak{c}} \mathfrak{D}^{\mathfrak{c}} \triangleleft \mathfrak{b}^{\mathfrak{b}} \mathbb{C} \triangleright \mathfrak{L} \mathfrak{L}^{\mathfrak{c}}$

[illegible]
$$(P = \langle b \rangle_{\mathcal{A}} \cap \mathcal{P} \cap \mathcal{A}^c, N = \langle b \rangle_{\mathcal{A}} \cap \mathcal{C} \cap \mathcal{A}^c \cap \mathcal{C}^c, M = \langle b \rangle_{\mathcal{A}} \cap \mathcal{C} \cap \mathcal{A}^c \cap \mathcal{C}^c, U = \langle b \rangle_{\mathcal{A}} \cap \mathcal{C} \cap \mathcal{A}^c \cap \mathcal{C}^c)$$

1. *Archives*. Grinnell also cites for Deane's collection.

- | | | |
|---|---------|---|
| 1 | polygon | Greiner Lake site for <i>Daphnia</i> collection |
| 2 | polygon | Cambridge Bay Sea-ice associated algae collection and copepod collection area |
| 3 | point | Outflow of sewage lagoon, leading into river/bay |
| 4 | point | Downstream of sewage lagoon, closest to outflow |
| 5 | point | Downstream of sewage lagoon, intermediate distance |
| 6 | point | Downstream of sewage lagoon, longest distance |
| 7 | point | Lake where Cambridge Bay drinking water is taken from |

